



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

IX. *On a new Property of the Tangents of three Arches trisecting the Circumference of a Circle*, by Nevil Maskelyne, D. D. F. R. S. and Astronomer Royal.

Read February 18, 1808.

MR. WILLIAM GARRARD having shewn me a curious property of the tangents of the three angles of a plane triangle, or in other words, of the tangents of three arches trisecting a semicircle, in a paper which I have communicated to this Society, I was led to consider whether a similar property might not belong to the tangents of three arches trisecting the whole circumference ; and, on examination, found it be so.

Let the circumference of a circle be divided any how into three arches A, B, C ; that is, let $A + B + C$ be equal to the whole circumference. I say, the square of the radius multiplied into the sum of the tangents of the three arches A, B, C, is equal to the product of the tangents multiplied together. I shall demonstrate this by symbolical calculation, now commonly called (especially by foreign mathematicians) analytic calculation.

It may be proper to premise, that the signification of the symbolical expressions of the tangents of an arc, whether with respect to geometry or numbers, are to be understood according to their position as lying on one side, or the other side of the radius, passing through the point of commencement of the arc of the circle ; those tangents which belong to

the first or third quadrant of the circle being considered as positive, and those belonging to the second and fourth quadrant, being of a contrary direction, as negative; in like manner as the sines in the first semi-circle are considered as positive, and in the second semi-circle as negative; and the cosines in the first and fourth quadrant are considered as positive, and in the second and third quadrants as negative; they lying, in the second case, on the contrary side of the diameter passing through the point of ninety degrees, to what they do in the former. Hence it easily follows, that the tangent of any arch and of its supplement to the whole circumference, or 360 degrees, are equal and contrary to one another, or the one negative of the other.

Let t, u, w , be put for the tangents of the three arches A, B, C respectively, and r for the radius, and \odot for the whole circumference. Then $A + B + C = \odot$, and $C = \odot - \overline{A + B}$.

By trigonometry, $t, \overline{A + B} = \frac{r^2 \times \overline{t + u}}{r^2 - tu}$, and the tang. C = tang.

$(\odot - \overline{A + B}) = -\text{tang. } \overline{A + B}$, by what has been said above.

Therefore $t, A + t, B + t, C$ or $t + u + w = t + u - \frac{r^2 \times \overline{t + u}}{r^2 - tu}$

$= tu \times -\frac{r^2 \times \overline{t + u}}{r^2 - tu}$; but t and u are the expressions for the tan-

gents of A and B respectively, and $-\frac{r^2 \times \overline{t + u}}{r^2 - tu}$ is the expression

for the tangent of C, or for w . Therefore, $r^2 \times \overline{t + u + w}$, or

the square of the radius multiplied into the sum of the three tangents of A, B, and C = tuw , or the product of the tangents.

Q. E. D.